



Review

Chromatographic analysis of vegetable oils: A review

Aluyor, E. O.^{1*}, Ozigagu, C. E.¹, Oboh, O. I.² and Aluyor, P.³

¹Department of Chemical Engineering, University of Benin, Benin City, Nigeria.

²Department of Chemical and Petroleum Engineering, University of Uyo, Uyo, Akwa Ibom State, Nigeria.

³Department of Vocation and Technical Education Ambrose Alli University, Ekpoma, Nigeria.

Accepted 10 March, 2009

Vegetable oils has become more attractive recently because of its economical benefits as they are used as components in many manufactured products and the fact that it is made from renewable resources. The determination of the minor components is of great importance in establishing the oil quality and their genuineness. The qualitative and quantitative determination of the constituents is done by gas chromatography of compounds or their derivatives. High precision in the gas chromatographic analysis of fatty acid methyl esters is possible with careful attention to detail during sample preparation, injection, chromatographic and data collection. This paper shows the applications of solid-phase extraction for the isolation of certain lipid classes prior to chromatographic analysis and also the most commonly used chromatographic techniques that can be applied for accuracy.

Key words: Chromatography, component, fatty acid methyl esters, sterol.

INTRODUCTION

Vegetable oils are mainly constituted by triacylglycerol (95 - 98%) and complex mixtures of minor compounds (2 - 5%) of a wide range of chemical nature. These minor constituents show a broad qualitative and quantitative composition depending on the vegetable species from which they are obtained. Agronomic and climatic conditions, fruit or seed quality, oil extraction system and refining procedures can cause variation in the content and composition of the constituents of vegetable oil (Cert et al., 2000).

The minor components include mono- and diglycerides, free fatty acids, phosphatides (or phospholipids), sterols, protein fragments, various resinous and mucilaginous materials and oxidative products (International Conference on Palm and Palm Products, 1989). A triglyceride is a chemical compound formed from one molecule of glycerol and three fatty acids.

Fatty acid chains may contain one or more double bonds at specific positions (unsaturated and polyunsaturated), or they may be fully saturated. The physical and

chemical properties of a fat depend on the composition of the fatty acid mixture. Fats from plant sources contain a higher proportion of unsaturated acids and are often liquids at room temperature due to hydrogen bonding. Polyunsaturated fats are usually of vegetable origin. Crisco is an example of a vegetable derived, unsaturated fatty acid that has been hydrogenated to form a solid material (Greenlief, 2004).

Fats are used in cooking because they are compounds with very high boiling points. Their high boiling points therefore make these classes of compounds ill suited for analysis by the gas chromatography. However, the glycerol ester can be chemically decomposed into methyl esters of each individual fatty acid (Greenlief, 2004). To analyze the minor component of vegetable oils, a preliminary qualitative and quantitative isolation step from the triacylglycerol matrix is required. Three basic procedures are normally used: saponification, liquid-liquid partition and chromatographic techniques. The saponification (heating with alcoholic solution of potassium hydroxide) transforms the glyceridic compounds in polar soaps allowing the extraction of unsaponified matters with hexane or diethyl ester.

Nevertheless, this procedure is not appropriate for wax

*Corresponding author. E-mail: eoaluyor@yahoo.com

esters, sterol esters, phenols, pigments, minor glyceridic compound and phospholipids, since they are altered during the saponification. Liquid-liquid partition with polar solvents (methanol-water, hexane-acetonitrile, etc.) is suitable for the isolation of phenols, polycyclic aromatic hydrocarbons and chlorophylls. Recently, supercritical fluids extraction has gained importance in separation techniques, due to the possibility of modifying product solubility through alteration of pressure and or temperature, or adding modifiers, substituting a wide variety of liquid solvent. Column chromatography is widely used to separate fractions having constituents of similar polarities. Actually, the latter is being substituted by solid-phase extraction (SPE), as it is a quicker technique and saves solvent volumes. The isolation of volatile compounds is achieved by gas stripping and distillation. The isolation fractions are still complex and require further fractioning, usually by means of thin-layer chromatography (TLC) or preparative high-performance liquid chromatography (HPLC) (Cert et al., 2000).

Gas chromatography (GC) and primarily high performance liquid chromatography (HPLC) are the two techniques widely applied for the analysis of major (triacylglycerols) and minor constituents of edible oils and fats (Christie, 1987; Cert et al., 2000). The qualitative and quantitative determination of the constituent is often done by capillary gas chromatography (GC) of the compounds or their derivatives. In general, GC assumes that the compound injected are volatile at the temperature of analysis and that they do not decompose at either the temperature of injection or analysis (Cert et al., 2000).

The history of the gas-liquid chromatograph of fatty acids dates back to 1952, when James et al. published a description of separations of free fatty acids from 1 to 12 carbon atoms. Later, they discovered that a previous methylation of fatty acids improved their volatility and their separation at temperatures below 250°C (James et al., 1956). A major advance was the introduction by Orr et al. (1958) of a liquid polyester phase as stationary phase able to conveniently separate saturated and unsaturated fatty acids. Later, the use of capillary columns with very films of various polar phases enabled to profoundly improve fatty acid analyses (Horning et al., 1963).

In the early days of GC, losses of esters of higher molecular weight were obtained because of chemical interactions with polyester liquid phases in packed columns. These problems were eliminated as catalytic-free polyesters available together with improve solid support materials. Perhaps the best objective assessment of the standard of fatty acid analysis possible with this older technology came in collaborative study of IUPAC method in 1979 (Firestone et al., 1979). For a range of fats and oils of commercial interest typical coefficients of variation (%) where quoted as 15 (2% level), 8.5 (5% level), 7 (10% level) and 3 (50% level).

Poorer data where obtain with difficult sample like milk

fat. In a comparable study with modern equipment, the results were only a little better (Beare-Rogers et al., 1990).

Preparation of fatty acids

The ability of an analytical method to characterize a vegetable oil is based on the identification and quantification of those compounds that are expected to be in connection with their origin and quality attributes (Cert et al., 2000).

Before analysis of vegetable oil samples can be commenced by any chromatographic procedure, it is first necessary to extract them from their tissue matrices in a relatively pure state (Christie, 1989). The analysis of fatty acids containing compound requires previously their hydrolysis or saponification, separation of the non-acidic constituent (when necessary) and the liberation of the acids from the mixture of soaps (James et al., 1963; Cert et al., 2000).

Adequate sample preparation techniques help to ensure an accurate and precise estimation of the true value of the food components.

Solid phase extraction (SPE)

Solid phase extraction (SPE) is a convenient approach for sample preparation in the Chromatographic analysis of edible fats as it requires a small amount of the test sample, low volumes of organic solvents, whereas the treatment is accomplished in a greatly shorter period of time than in other techniques.

SPE typically involves commercially available pre-packed polymer or glass mini disposable columns (cartridges) or disks that can be used to remove contaminants or to further fractionate the analyses prior to analysis. The technique is highly selective and versatile due to the large number of phases introduced in the last ten years. New formats (96-well plates, micro fibers for solid phase micro extraction), automation and on-line use are the important advantages for the current and future applications of SPE.

APPLICATION OF CHROMATOGRAPHIC ANALYSIS ON VEGETABLE OILS

Sterols: Sterols free are esterified with fatty acids, and are found in small concentrations (0.1 - 5.6%) in edible fats. Sterols, such as cholesterols are alcohols with the cyclopentanophenanthrene ring system (atom 1 through 17 as in Figure 1). This substructure is also found in steroid hormones such as testosterone, progesterone, and cortisol. Cholesterol is found in animal fats and at relatively low amounts in certain plant lipids (e.g. palm oil). Cholesterol is classified as an alcohol because it has a




Figure 1. Structural diagrams of some Sterols. (Zamora, 2005).

Table 1. Extraction of sterol fraction according to Toivo et al., (1998)

Step	Action
A	Saponification of sample [0.25-0.50g oil + 5ml 0.5 mM ethanolic KOH, 20min reflux, 80°C]
B	Addition of betulin as internal standard [1 mg betulin in 5ml CHCl ₃]
C	PH adjustment, filtration [few drops 5M HCl, 0.45 mm nylon membrane filter]
D	Application to C ₁₈ cartridge

a hydroxyl group (-OH) in position 3 of the ring system. Cholesterol is produced by the liver and is found in all body tissues where it helps to organize cell membranes and control their permeability. Cholesterol derivatives in the skin are converted to vitamin D when the skin is exposed to sunlight. Vitamin D₃ mediates intestinal calcium absorption and bone calcium metabolism. A high level of cholesterol in the blood is considered to be a risk factor for cardiovascular diseases. Cholesterol and lipoprotein levels can be normalized through exercise and reduced calorie diets that eliminate hydrogenated fats and add sources of polyunsaturated fatty acids such as grape oil (Zamora, 2005).

Phytosterols, the plant sterols, are rather simple mixtures of closely related organic compounds. Some of

them always accompany each other (e.g. β -sitosterol and campesterol). The analysis of sterols is important for detecting oil adulteration, e.g. of butter fat with vegetable oils or of virgin olive oil with cheaper oils. More recent is the interest in the nutritional value of sterols and the monitoring and accurate quantification of oxysterols. The latter, especially cholesterol oxidation products, have been examined thoroughly due to their potential health risk (Bosinger et al., 1993).

Capillary Gas Chromatography (CGC) is mainly used for the analysis of sterols. The sterol fraction is obtained from the unsaponified matter by Thin Layer Chromatography (TLC) prior to analysis. This tedious procedure, part of the official ones adopted for the control of edible oils (EC, 1991; IURJ/C, 1992; AOAC, 1999) has a little potential for automation and limits the number of samples that can be handled daily. Alternative sample preparation procedures involve the derivation of the unsaponified (Slover et al., 1983), automatic removal of the interfering triacylglycerols by continuous on-line trans-esterification with potassium methylate in methanol (Ballesteros et al., 1996), fractionation on Column Chromatography (CC) (Homborg, 1987) or on-line LC-GC (Grob et al., 1989). Using these alternative procedures, tocopherols, hydrocarbons and related compounds are also determined in the subsequent analysis. A procedure was applied to crude, extra virgin, refined and crude pomace olive oils. Toivo et al. (1998) extended the efforts made by Tsui (1989) on C₁₈ packing material to clean up the unsaponifiables according to the scheme presented in Table 1 above. The merits of HPLC procedures have also been investigated in the literature for their potential, either for sample treatment, as direct analytical means or in conjunction with other chromatographic techniques (Holen, 1985; Cert et al., 2000). Thus, there is an increasing interest in developing rapid methods for sterol analysis (Alonso et al., 1997). SPE procedures in the analysis of sterols in edible fats are rather limited though more information can be found in the case of biological fluids.

In all methods 500 mg of milk fat spiked with 30 mg of COPs was applied to the columns. A combination of silica SPE and NH₂ cartridges given in detail in Table 2 (below) allowed the optimum removal of interfering components and the reliable separation (free of spurious peaks) and quantification of COPs.

Wax esters: The wax ester formed by the reaction of alcohols (aliphatic, triterpenic, methylsterols and sterols) and free fatty acids are present in seed and fruit. During the oil extraction process, a fraction of these esters is transferred into the oil depending on the oil extraction system. So solvent extracted oils contain higher concentration of wax esters compared with cold-pressed and centrifuged ones. C₄₀, C₄₂, C₄₄ and C₄₆ waxes, deriving from straight chain alcohols are very abundant in olive pomace oils while low concentrations are found in olive

Table 2. Combined elution of COPs using silica and NH₂ SPE columns (Lai et al. 1995; Ulberth and Rössler 1998)

Step	Action
A	LC-Si SPE column (300mg/3ml); Conditioning with 3 ml hexane
B	Sample application
C	Elution with 10 ml n-hexane/ethyl ether (95:5,v/v); 25 ml n-hexane/ethyl ether [90 10.15 ml n-hexane/ethyl ether (80:20. v/v), vacuum 29 kPa, solvent flow rate 0.6 ml/mm v/v]
D	Elution of COPs with 5 ml acetone
E	Concentration of the eluate to 1 ml of n-hexane/ethyl ether (90:10, v/v)
F	Application to NH ₂ -SPE column (500 mg)
G	Elution with 15 ml of n-hexane/ethyl ether (90:10, v/v)
H	Elution of purified COPs with 10 ml acetone

oils. In addition, great concentration of waxes yields turbidness during refining or storage. Consequently, the determination of waxes is important to evaluate the quality and genuineness of some vegetable oils (Cert et al., 2000).

Other alcohols and related derivatives: Waxes are esters of long chain alcohols with fatty acids. They are found in vegetable oils and their levels depend on the technology of extraction from fruits and seeds. The analysis of waxes is important in the quality control of de-waxing and winterization processes (Mariani et al., 1987). The composition of waxes varies in the various oils so that information can be obtained for the presence of inferior quality oils as it is established in the case of virgin olive oil (EC, 1993). The wax esters are obtained from the oil after saponification and then analyzed by capillary GC. Simplification of the sample preparation step has been attempted by several authors (Grob et al., 1989 and 1990; Amelio et al., 1993). Reiter et al. (1999), who applied for the first time SPE to isolate waxes from fennel and caraway seed oils, claimed that the procedure is applicable to other vegetable oils. The oil (30 mg / 500 μ l n-hexane) containing the internal standard was applied to silica cartridges (500 mg / 3 ml, Bakerbond, J.T. Baker) preconditioned with the same solvent. The sample was drawn through the column at a flow rate of 2.5 ml/min (controlled by a vacuum manifold processor) and the waxes were then eluted with n-hexane: diethyl ether (99:1, v/v) (10 ml). Recovery studies were very satisfactory. Yields from the capillary gas chromatographic analysis for stearic acid palmityl ester and behenic acid behenyl ester (40 mg/ml, each) ranged between 95 - 98% (SD % < 5). Nota et al. (1999) simplified the method of isolation of waxes of olive oil in an attempt to improve the existing official method for their determination. The separation was carried out on SPE cartridges (1 g Chromabond, Macherey-Nagel) rinsed first with carbon tetrachloride (6 ml). Oil (20 mg) plus the internal standard was

transferred onto the column with a minimum amount of carbon tetrachloride (3 x 100 μ l) and the waxes were eluted with the same solvent (6 ml). The speed of elution ranged between 0.5 - 1.5 ml/min. High levels of free acidity (up to 10%) did not lower the accuracy of the method.

Phenolic compounds: Phenolic compounds are transferred from fruits and seeds to raw oils during processing. Refining eliminates the content of phenolic compounds; therefore, their presence is mainly of interest in virgin olive oil and to a certain extent in cold-pressed seed oils, e.g. cold -pressed canola oil. The importance of phenolic compounds in the sensory quality and the stability of virgin olive oil and their effect to human health has been discussed (Tsimidou, 1998). The major phenolics found in virgin olive oil are hydroxytyrosol, tyrosol, aldehydic secoiridoids, flavonoids and lignans (acetoxypinoresinol, pinoresinol) (Brenes et al., 2000). The content of these natural antioxidants depends on the olive oil processing (two-phase extractors yield in general oils with more phenols) and storage conditions. Except from the colorimetric procedures using the Folin-Ciocalteu reagent for the determination of total phenols, gas chromatographic methods and, basically, HPLC procedures have been used for the analysis of the oil fraction containing the phenolic compounds (Ryan and Robards, 1998).

Liquid-liquid extraction and more recently solid-phase extraction have been used for the isolation of the so-called "polar fraction". The solvent system usually applied is aqueous methanol in various proportions. Tsimidou (1999) has recently reviewed the advances in extraction and analysis of virgin olive oil polyphenols. Perez-Camino (2000) presented also applications of solid -phase extraction of phenolic compounds of virgin olive oil. The first systematic study was carried out by Papadopoulos and Tsimidou (1992), who, in a comparative study of two SPE systems and a liquid-liquid one, found, using HPLC, that polyvinylpyrrolidone (PVP) traps selectively the phenolic

compounds. C_{18} cartridges of the same bed volume and with the same solvent system did not retain the bound forms of phenolic compounds. Recovery studies using hydroxytyrosol and tyrosol on an oil matrix (40 mg/kg each) gave yields of 50 and 90%, respectively. In 2001, Liberatore et al., observed the non-homogeneous and even contradictory results concerning the recovery of phenolics using different methods. They reported recovery data for eight representative olive oil phenols (tyrosol, p-hydroxybenzoic, vanillic, protocatechuic, syringic, p-coumaric, ferulic and caffeic acids) at two levels of addition (50 and 500 mg/kg) using three different extraction procedures. The efficiency of two SPE procedures (C_{18} and C_8 end capped) was compared with that of a liquid-liquid extraction previously reported by Tsimidou et al. (1992). End capped reversed-phase was proved less efficient than C_{18} . Suppression of the residual polar groups of silica had an adverse effect on the release mechanism of the analytes. Still, the recovery data were not consistent at the two levels of addition to permit a better understanding of retention of components of similar molecular weight and polarity.

Partial glycerides and free fatty acids: The analysis of partial glycerides, mono- and di-glycerides is usually carried out by capillary gas chromatography. HPLC may only give information on the total amounts of different classes of partial glycerides (e.g. 1, 2- and 1, 3- di-glycerides). These compounds, intermediates in the biosynthesis of triglycerides, are also markers of oil freshness or of employed technological treatments. Perez-Camino et al., (1996) reported on alterations on di-glyceride composition using different types of SPE columns (di-ol, silica, amino). They found that only the di-ol phase did not cause isomerization during the extraction. In their recent review (Perez-Camino, 2000), the authors commenting on these and other literature data stated that "a lot of care must be taken before choosing a support until more explanation and importance is given to the absorbent employed". Similar observations were also made by Conte et al. (1997).

Free fatty acid content is not a problem in the analysis of vegetable fats that are consumed refined with the exception of virgin olive oil. Hopia et al. (1992), describing the analysis of lipid classes by solid-phase extraction gave information for the elution of free fatty acids from edible oils and fats. Procedures related to this group of lipids is found for other categories of food products, where the level and composition of free fatty acids, in particular short chain ones, affect the organoleptic characteristics of the finished products (e.g. cheese) (Innocente et al., 2000).

Phospholipids / Phosphatides: Phospholipids or phosphatides are natural surfactants and emulsifiers consisting of an alcohol such as glycerol, one or two molecules

of fatty acids and a phosphoric acid compound. They are found in all plants and animals and include such substances as lecithin, cephalin, and sphingomyelin. Lecithin is a significant constituent of brain and nervous tissue consisting of a mixture of diglycerides of stearic, palmitic, and oleic acid linked to the choline ester of phosphoric acid (Zamora, 2005).

Phospholipids contribute to the stability and quality of edible oils, fats and fatty foods through their antioxidative activity or contribution to the texture (Singleton, 1993). On the other hand, they are responsible for oil discoloration during deodorization and steam distillation so that their determination is necessary to evaluate the efficiency of degumming (Mounts and Nash, 1990). Some oils contain substantial amounts of phospholipids, e.g. soybean oil, sunflower oil and canola oil or peanut oil (Przybyski and Eskin, 1991). The most important members of this class of lipids found in edible fats are phosphatidylethanolamine, phosphatidylinositol, phosphatidylserine and phosphatidyl choline. Egg yolk and milk fat also contain significant amount of sphingomyelin. The fatty acid composition of the various phospholipids depends on the fat origin (Lendrath et al., 1991). Fish oils contain similar mammalian oil phospholipids. After refining oils and fats of vegetable origin have rather low amounts so that a concentration step is needed prior to HPLC, which is the preferred method that replaced thin layer chromatography (Singleton, 1993).

SPE has been employed in certain cases for the pre-concentration of phospholipids. Nash and Frankel (1986) fractionated degummed, crude soybean oils using silicic acid columns (Sep-Pak, Waters Assoc.) prior to HPLC analysis. Oil, 1 g of a 10% solution in petroleum ethyl ether (95:5, v/v), was applied, the non-polar lipids were eluted with the same system (10 ml), the intermediate polarity lipids were eluted with diethyl ether (20 ml) and finally the phospholipids were obtained with methanol (10 ml). The applicability of the system was checked by thin layer chromatography of the various fractions. Caboni et al., (1996) separated phospholipids from total lipids that had been extracted from food samples (egg powder, chicken meat, ripened cheese and salami) by the Folch method. The lipid extract was purified by using normal phase (silica) ion exchange (aminopropyl) and reversed-phase (C_{18} and C_8) SPE cartridges (500 mg, BondElut, Varian). The purified fractions were analyzed by HPLC coupled with an evaporative light-scattering detector. Details of the four SPE elution systems are given in Table 3.

Based on phosphorous determination in the various fractions, the authors reported that phospholipids eluted in the third fraction from polar phases (silica and amino) and in the first fraction derived from the reversed-phase bonded materials. Though precision data were satisfactory in all four procedures, accuracy data varied significantly. C_8 packing was proved more efficient for individual and

Table 3. SPE Separation of phospholipids from fatty foods according to Caboni et al. (1996)

SPE mode *	Lipid sample	Conditioning solvent system	Elution system (v/v/v)
Silica	200mg/0.5ml CHCl ₃	3 ml n-hexane	4 ml n-hexane:diethyl ether (8:2); 4 ml n-hexane:diethyl ether (1:1); 4 ml methanol or 4ml 1-5% CH ₃ COOH in methanol
Amino-propyl	200mg/0.5ml CHCl ₃	3 ml n-hexane	2x2.5 ml CHCl ₃ :isopropanol (2:1); 2x2.5 ml 2% CH ₃ COOH in diethyl ether; 4x1ml methanol
C ₁₈	200mg/0.5ml CHC I ₃ :Methanol (2:1, v/v)	3m/methanol	4ml methanol/water (4:1); 4ml methanol; 4ml methanol; CHCl ₃ (4:1);
C ₈	200gm/0.5ml CHCl ₃ : methanol (2:1, v/v)	3ml/methanol	4 ml methanol; 5 ml CHCl ₃ : methanol (3:2) 5 ml CHCl ₃

*Column regeneration with 3 ml CHCb; 3 ml CHCl₃: methanol (2:1); 5 ml methanol.

total phospholipid recoveries. Carelli et al. (1997) enriched sunflower oil samples with phospholipids using SPE cartridges and subsequently performed HPLC analysis based on the IUPAC standard method for soybean lecithin and ultraviolet detection. The bonded di-ol cartridges (500 mg, J. T. Baker) were preconditioned with methanol (2 ml), chloroform (2 ml) and hexane (4 ml). From the oil dissolved in chloroform (50 - 150 mg) triglycerides were removed with chloroform (2.5 ml). Phospholipids were eluted with 7 ml methanol that contained 0.5 ml / 100 ml of a 25% ammonia solution. Accuracy and precision data reported were quite satisfactory. Phosphatidylcholine, phosphatidylethanolamine, phosphatidylserine and phosphatidylinositol were sequentially eluted from amino-propyl bonded silica with acetonitrile/n-propanol (2:1, v/v), methanol, isopropanol/methanolic HCl (4:1, v/v) and methanol/methanolic HCl (9:1, v/v). The recoveries were excellent (Pietsch and Lorenz, 1993).

Conclusion

The qualitative and quantitative determination of the major and minor constituents of vegetable oils are done by gas chromatography and primarily high performance liquid chromatography which are the two techniques widely applied for the analysis of edible oils and fats.

Solid phase extraction (SPE) is a simple approach for sample preparation in the Chromatographic analysis of edible fats as it requires small amounts of the test samples, low volumes of organic solvents, whereas the treatment is accomplished in a greatly shorter period of time than in other techniques. In the course of this work the merits of HPLC procedures have been shown for their potential, either for sample treatment, as direct analytical means or in conjunction with other chromatographic tech-

iques.

REFERENCES

- Alonso L, Fontecha J, Lozada L, Juarez M (1997). Determination of mixtures of vegetable oils and milk fat by analysis of sterol fraction by gas chromatography. *J. Am. Oil Chem. Soc.* 74: 131-135.
- Amelio M, Rizzo R, Varazini F (1993). Separation of wax esters from olive oils by high-performance liquid chromatography. *J. Am. Oil Chem. Soc.* 70: 793-796.
- Ballesteros E, Gallego M, Vajcarcel M (1996). Gas chromatographic determination of cholesterol and tocopherols in edible oils and fats with automatic removal of interfering triglycerides. *J. Chromatogr. A* 719: 221-227.
- Bosinger S, Luf W, Braridl E (1993). Oxysterols: (heir occurrence and biological effects. *Int. Dairy Sci.* 3: 1-33.
- Brenes M, Hidalgo FJ, Garcia A, Rios JJ, Garcia P, Zamora R, Garrido A (2000). Pinoresinol and 1-acetoxypinoresinol, two new phenolic compounds identified in olive oil. *J. Am. Oil Chem. Soc.* 77: 715-720.
- Caboni MA, Menotta S, Lercker G (1996). Separation and analysis of phospholipids in different foods with a light-scattering detector. *J. Am. Oil Chem. Soc.* 73: 1561-1566.
- Cert A, Moreda W, Perez C (2000). Chromatographic analysis of minor constituents in vegetable oils. *J. Chromatogr. A* 881: 131 -148.
- CH 4200, Fall 2004, Prof. Greenlief.
- Christie WW(1989). Preparation of ester derivatives of fatty acids for chromatographic analysis. In: *Advances in Lipid Methodology – Two.* (Ed . W. W. Christie, Oily Press, Dundee) pp. 69-111.
- Conte LS, Koprivnjak O, Fiorasi S, Pizzale L (1997). Applicazione della estrazione in fase solida (SPE) alia determinazione dei diacilgliceroli negli alimenti. *Riv. H. Sost. Grasse* 74: 411-414.
- EC Regulation 183/29.1.93. (1993). Official Journal of the European Communities. Modifying of the EC Regulation 2568/91 regarding the characteristics of olive oils and kernel oils and their methods of analysis L.248.
- EC Regulation 2568/1.7.91. (1991). Official Journal of the European Communities. On the characteristics of olive oils and kernel olive oils and on their methods of analysis L248.
- Firestone D, Horowitz W (1979) IUPAC gas chromatography method for determination of fatty acid composition: collaborative study. *J. Assoc. Off. Anal. Chem.* 62: 709-721.
- Grob K, Lanfranchi C, Mariani C (1990). Evaluation of olive oils through

- the fatty alcohols, the sterols and their esters by coupled LC-GC. *J. Am. Oil Chem. Soc.* 67: 626-634.
- Holen B (1985). Rapid separation of free sterols by reversed-phase HPLC. • « fa n*rt*(* W^O. - *J. Am. Oil Chem. Soc.* 62: 1344-1346.
- Homberg E (1987). Rapid method for the isolation of sterols from fats and oils. *Fat Sci. Technol. Q.* pp. 215-216.
- Hopia AI, Piironen VI, Koivistoinen PE, Hyvonen LET (1992). Analysis of lipid classes by solid-phase extraction and high-performance size-exclusion chromatography. *J. Am. Oil Chem. Soc.* 69: 772-776.
- Innocente N, Moret S, Corradini C, Conte LS (2000). A rapid method for the quantitative determination of short chain free volatile fatty acids from cheese. *J. Agric. Food Chem.* 48: 3321-3323.
- Lai SM, Gray JI, Zabik ME (1995). Evaluation of solid-phase extraction and gas chromatography for the determination of cholesterol oxidation products in spray-dried whole egg. *J. Agric. Food Chem.* 43: 1122-1126.
- Lendrath G, Bonekamp A, Kraus Lj (1991). Quantitative planar chromatography of phospholipids with different fatty acid compositions. *J. Chromatogr.* 588: 303-305.
- Mariani C, Lanzani A, Fedeli E (1987). La presentiadi cere nell'olio di girasole in funzione di alcuni trattamenti tecnologici. *Riv.It. Sost. GrasseLXIV.* pp.13-16.
- Mounts TL, Nash AM (1990). HPLC analysis of phospholipids in crude oil for evaluation of soybean deterioration. *J. Am. Oil Chem. Soc.* 67: 757-760.
- Nash AM, Frankel EN (1986). Limited extraction of soybeans with hexane. *J. Am. Oil Chem. Soc.* 63: 244-246.
- Nota G, Spagna MS, Romano R, Naviglio D, Impronta C (1999). Idrolisi rapida degli esteri degli steroli nei grass!. *Riv. It. Sost. Grasse.* 72: 315-316.
- Pietsch A, Lorenz RL (1993). Rapid separation of major phospholipid classes on a single aminopropyl cartridge. *Lipids2S,* pp. 945-947.
- Przybyski R, Eskin NAM (1991). Phospholipid composition of canola oils during the early stages of processing as measured by TLC with flame ionization detector. *J. Am. Oil Chem. Soc.* 68: 241 -245.
- Reiter B, Lechner M, Lorbeer E, Aicholz R (1999). Isolation and characterization of wax esters in fennel and caraway seed oils by SPE-GC. *J. High Resol. Chromatogr.* 22: 514-520.
- Ryan D, Robards K (1998). Phenolic compounds in olives. *Analyst,* 123: 31R-44R.
- Toivo J, Piironen V, Kalo R, Varo P (1998). Gas chromatographic determination of major sterols in edible oils and fats using solid-phase extraction in sample preparation. *Chromatographia* 48: 745-750.
- Tsimidou M (1998). Polyphenols and Quality of Virgin Olive Oil in Retrospect. *Ital. J. Food Sci.* 10: 99-116.
- Tsui IC (1989). Rapid method of total cholesterol in homogenized milk. *J. Assoc. Off. Anal. Chem.* 72: 421-424.
- Ulberth R, Rossler F (1998). Comparison of solid-phase extraction methods for the clean up of cholesterol oxidation products. *J. Agric. Food Chem.* 46: 2634-2637.
- Zamora Antonio (2005), *Fats, Oils, Fatty Acids, Triglycerides.* Scientific Psychil.